



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b>  <b>A61K 31/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <span style="float: right;"><b>WO 00/12074</b></span>  <b>(43) International Publication Date:</b> <span style="float: right;">9 March 2000 (09.03.00)</span>		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top; padding: 5px;"> <b>(21) International Application Number:</b> PCT/US99/19845   <b>(22) International Filing Date:</b> 27 August 1999 (27.08.99)   <b>(30) Priority Data:</b>  <div style="display: flex; justify-content: space-between;"> <div>60/098,219</div> <div>28 August 1998 (28.08.98)</div> <div>US</div> </div> <div style="display: flex; justify-content: space-between;"> <div>60/125,343</div> <div>19 March 1999 (19.03.99)</div> <div>US</div> </div>   <b>(71) Applicant (for all designated States except US):</b> SCIOS INC.  [US/US]; 820 West Maude Avenue, Sunnyvale, CA 94086 (US).   <b>(72) Inventors; and</b>  <b>(75) Inventors/Applicants (for US only):</b> GOEHRING, R., Richard [US/US]; 6195 Potters Lane, Pipersville, PA 18947 (US). LUEDTKE, Gregory, R. [US/US]; Apartment B, 963 La Meso Terrace, Sunnyvale, CA 94086 (US). MAVUNKEL, Babu, J. [US/US]; 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). CHAKRAVARTY, Sarvajit [IN/US]; 976-2 Alpine Terrace, Sunnyvale, CA 94086 (US). DUGAR, Sundeep [IN/US]; 749 Wingate Drive, Bridgewater, NJ 08807 (US). SCHREINER, George, F. [US/US]; 12274 Leander Drive, Los Altos Hills, CA 94022 (US). LIU, David, Y. [US/US]; 201 Ferne Avenue, Palo Alto, CA 94036 (US). LEWICKI, John, A. [US/US]; 308 Escobar Avenue, Los Gatos, CA 95030 (US). </td> <td style="width: 50%; vertical-align: top; padding: 5px;"> <b>(74) Agents:</b> FAVORITO, Carolyn et al.; Morrison &amp; Foerster LLP, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).   <b>(81) Designated States:</b> AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, EE, GE, HU, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).   <b>Published</b>  <i>Without international search report and to be republished upon receipt of that report.</i> </td> </tr> </table>			<b>(21) International Application Number:</b> PCT/US99/19845  <b>(22) International Filing Date:</b> 27 August 1999 (27.08.99)  <b>(30) Priority Data:</b> <div style="display: flex; justify-content: space-between;"> <div>60/098,219</div> <div>28 August 1998 (28.08.98)</div> <div>US</div> </div> <div style="display: flex; justify-content: space-between;"> <div>60/125,343</div> <div>19 March 1999 (19.03.99)</div> <div>US</div> </div> <b>(71) Applicant (for all designated States except US):</b> SCIOS INC. [US/US]; 820 West Maude Avenue, Sunnyvale, CA 94086 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GOEHRING, R., Richard [US/US]; 6195 Potters Lane, Pipersville, PA 18947 (US). LUEDTKE, Gregory, R. [US/US]; Apartment B, 963 La Meso Terrace, Sunnyvale, CA 94086 (US). MAVUNKEL, Babu, J. [US/US]; 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). CHAKRAVARTY, Sarvajit [IN/US]; 976-2 Alpine Terrace, Sunnyvale, CA 94086 (US). DUGAR, Sundeep [IN/US]; 749 Wingate Drive, Bridgewater, NJ 08807 (US). SCHREINER, George, F. [US/US]; 12274 Leander Drive, Los Altos Hills, CA 94022 (US). LIU, David, Y. [US/US]; 201 Ferne Avenue, Palo Alto, CA 94036 (US). LEWICKI, John, A. [US/US]; 308 Escobar Avenue, Los Gatos, CA 95030 (US).	<b>(74) Agents:</b> FAVORITO, Carolyn et al.; Morrison & Foerster LLP, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).  <b>(81) Designated States:</b> AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, EE, GE, HU, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(21) International Application Number:</b> PCT/US99/19845  <b>(22) International Filing Date:</b> 27 August 1999 (27.08.99)  <b>(30) Priority Data:</b> <div style="display: flex; justify-content: space-between;"> <div>60/098,219</div> <div>28 August 1998 (28.08.98)</div> <div>US</div> </div> <div style="display: flex; justify-content: space-between;"> <div>60/125,343</div> <div>19 March 1999 (19.03.99)</div> <div>US</div> </div> <b>(71) Applicant (for all designated States except US):</b> SCIOS INC. [US/US]; 820 West Maude Avenue, Sunnyvale, CA 94086 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GOEHRING, R., Richard [US/US]; 6195 Potters Lane, Pipersville, PA 18947 (US). LUEDTKE, Gregory, R. [US/US]; Apartment B, 963 La Meso Terrace, Sunnyvale, CA 94086 (US). MAVUNKEL, Babu, J. [US/US]; 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). CHAKRAVARTY, Sarvajit [IN/US]; 976-2 Alpine Terrace, Sunnyvale, CA 94086 (US). DUGAR, Sundeep [IN/US]; 749 Wingate Drive, Bridgewater, NJ 08807 (US). SCHREINER, George, F. [US/US]; 12274 Leander Drive, Los Altos Hills, CA 94022 (US). LIU, David, Y. [US/US]; 201 Ferne Avenue, Palo Alto, CA 94036 (US). LEWICKI, John, A. [US/US]; 308 Escobar Avenue, Los Gatos, CA 95030 (US).	<b>(74) Agents:</b> FAVORITO, Carolyn et al.; Morrison & Foerster LLP, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).  <b>(81) Designated States:</b> AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, EE, GE, HU, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>			
<b>(54) Title:</b> INHIBITORS OF p38- $\alpha$ KINASE				
<div style="text-align: right; margin-top: -20px;">(1)</div>				
<b>(57) Abstract</b>  <p>The invention is directed to methods to treat conditions mediated by kinase using compounds of formula (1) and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof, wherein Z is N or CR<sup>1</sup>, R<sup>1</sup> is a noninterfering substituent, each of X<sup>1</sup> and X<sup>2</sup> is a linker, Ar<sup>1</sup> and Ar<sup>2</sup> are identical or different, and represent optionally substituted C1-C20 hydrocarbyl residues wherein at least one of Ar<sup>1</sup> and Ar<sup>2</sup> is an optionally substituted aryl group, with the proviso that when X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof, X<sup>1</sup> is CO or an isostere thereof, and Ar<sup>2</sup> is optionally substituted phenyl, Ar<sup>1</sup> is other than an optionally substituted indolyl, benzimidazolyl or benzotriazolyl substituent, and wherein said optionally substituted phenyl is not an optionally substituted indolyl, benzimidazolyl, or benzotriazolyl, Y is a noninterfering substituent, wherein n is an integer from 0-4, and wherein m is an integer from 0-4 and l is an integer from 0-3.</p>				

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## INHIBITORS OF p38- $\alpha$ KINASE

### Technical Field

The invention is directed to compounds that are useful in treating inflammation and  
5 that contain N-containing heterocycles such as piperazine or piperidine moieties coupled  
to phenyl and other aryl groups. More particularly, the invention concerns novel  
compounds of this type as well as methods to treat heart and kidney conditions using these  
and other compounds.

### 10 Background Art

A large number of chronic and acute conditions have been recognized to be  
associated with perturbation of the inflammatory response. A large number of cytokines  
participate in this response, including IL-1, IL-6, IL-8 and TNF. It appears that the  
activity of these cytokines in the regulation of inflammation rely at least in part on the  
15 activation of an enzyme on the cell signaling pathway, a member of the MAP kinase  
family generally known as p38 and alternatively known as CSBP and RK. This kinase is  
activated by dual phosphorylation after stimulation by physiochemical stress, treatment  
with lipopolysaccharides or with proinflammatory cytokines such as IL-1 and TNF.  
Therefore, inhibitors of the kinase activity of p38 are useful antiinflammatory agents.

20 PCT applications WO98/06715, WO98/07425, WO98/28292 and WO 96/40143,  
all of which are incorporated herein by reference, describe the relationship of p38 kinase  
inhibitors with various disease states. As mentioned in these applications, inhibitors of  
p38 kinase are useful in treating a variety of diseases associated with chronic  
inflammation. These applications list rheumatoid arthritis, rheumatoid spondylitis,  
25 osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic  
shock, Gram-negative sepsis, toxic shock syndrome, asthma, adult respiratory distress  
syndrome, stroke, reperfusion injury, CNS injuries such as neural trauma and ischemia,  
psoriasis, restenosis, cerebral malaria, chronic pulmonary inflammatory disease, silicosis,  
pulmonary sarcosis, bone resorption diseases such as osteoporosis, graft-versus-host  
30 reaction, Crohn's Disease, ulcerative colitis including inflammatory bowel disease (IBD)  
and pyresis.

The above-referenced PCT applications disclose compounds which are p38 kinase inhibitors said to be useful in treating these disease states. These compounds are either imidazoles or are indoles substituted at the 3- or 4-position with a piperazine or piperidine ring linked through a carboxamide linkage. Additional compounds which are conjugates of piperazines with indoles are described as insecticides in WO97/26252, also incorporated herein by reference.

U.S. Patent No. 5,719,135 describes tyrosine kinase inhibitors containing piperidine or piperazine rings linked through a methylene at position 1 of piperidine to various aromatic systems which must further contain a  $\gamma$  lactam fused to a pyridine ring.

10 Similar compounds are described in U.S. Patent No. 5,663,346 and in WO96/22976.


Other cyclic tyrosine kinase inhibitors are described in PCT application WO95/06032. In addition, WO94/20062 describes balanoids as protein kinase C inhibitors. The balanoid compounds contain multiple aromatic systems which include at least a ring containing at least seven members. Some of the compounds useful in the method of the present

15 invention are known compounds.

## Disclosure of the Invention

The invention is directed to methods of treating inflammation generally, including specific conditions such as those described in the Background section above. The compounds of the invention have been found to inhibit p38 kinase and are thus useful in treating diseases mediated by this enzyme. These compounds also inhibit p38 $\alpha$  preferentially as compared to their inhibition of p38 $\beta$  as is further discussed below.

The compounds useful in the invention are of the formula



25                    and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof,

wherein


Z is N or CR<sup>1</sup>, R<sup>1</sup> is a noninterfering substituent,

each of X<sup>1</sup> and X<sup>2</sup> is a linker,

Ar<sup>1</sup> and Ar<sup>2</sup> are identical or different, and represent optionally substituted C1-C20 hydrocarbyl residues wherein at least one of Ar<sup>1</sup> and Ar<sup>2</sup> is an optionally substituted aryl group, with the proviso that when X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof, X<sup>1</sup> is CO or an isostere thereof, and Ar<sup>2</sup> is optionally substituted phenyl, Ar<sup>1</sup> is other than an optionally substituted indolyl, benzimidazolyl or benzotriazolyl substituent, and wherein said optionally substituted phenyl is not an optionally substituted indolyl, benzimidazolyl, or benzotriazolyl,

Y is a noninterfering substituent, wherein n is an integer from 0-4, and wherein m is an integer from 0-4 and l is an integer from 0-3.

Preferably, the compounds useful in the invention are of the formula



and the pharmaceutically acceptable salts thereof,

wherein Ar<sup>1</sup> is optionally substituted furanyl, thiophenyl, phenyl system having 0, 1, or 2 heterocyclic N atoms or naphthyl system having 0, 1, 2 or 3 heterocyclic N atoms;

X<sup>1</sup> is CO or an isostere thereof;

Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C);

n is 0 or 1;

Z is CH or N;

X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof; and

Ph is optionally substituted phenyl.

The optional substituents on the aryl moieties (including phenyl) include halo, nitro, optionally substituted alkyl (1-6C) or alkenyl (1-6C), CN, guanidino or CF<sub>3</sub>, as well as RCO, COOR, CONR<sub>2</sub>, SO<sub>2</sub>NR<sub>2</sub>, -OOCR, -NROCR, -NROCOR, NR<sub>2</sub>, OR or SR, wherein R is H or alkyl (1-6C), as well as substitution by phenyl, itself optionally

substituted by the foregoing substituents. Any two substituents may form a 5-7 membered carbocyclic or heterocyclic ring subject to the proviso.

Thus, in one aspect, the invention is directed to compounds of the formulas set forth above and to pharmaceutical compositions containing them. In other aspects, the invention is directed to methods of treatment using compounds of the formula set forth  
5 above. The invention is also directed to specific classes of compounds within the genus of formula (1). In other aspects, the invention is directed to methods to produce the classes of compounds useful in the invention.

#### 10 Modes of Carrying Out the Invention

The compounds of formula (1) set forth above are defined by the nature of the substituents on the heterocycloalkyl ring in the center of the formula. Piperazine or piperidine rings are preferred and piperidine rings are more preferred. The central piperazine or piperidine ring can be expanded or contracted using -CH<sub>2</sub> groups so that it  
15 includes from 4 members up to 11 members. The substitution on this ring is on the N or Z positions only. Although not bound by this theory, the function of the central heterocycloalkyl group is apparently to space the Ar<sup>2</sup> group, which is generally hydrophobic, from the Ar<sup>1</sup> group, which is preferably but not necessarily hydrophilic.

The central heterocycloalkyl ring can include from 1-2 N. If the ring contains only  
20 1 N then Z is a -CR<sup>1</sup> wherein R<sup>1</sup> is a noninterfering substituent. Preferably, R<sup>1</sup> is alkyl, alkoxy, aryl, arylalkyl, aryloxy, heteroaryl, halogen, acyl, carboxy, or hydroxy. More preferably, R<sup>1</sup> is hydroxy, alkyl, or alkoxy.

The linker that couples the central heterocycloalkyl ring and the Ar<sup>1</sup> and Ar<sup>2</sup> groups on either end of the molecule are preferably saturated or unsaturated alkylene optionally  
25 containing 1-4 carbonyl, 1-4 SO<sub>2</sub>, and/or 1-3 heteroatoms, including a linker which is CO, SO<sub>2</sub>, SO or contains a heteroatom, and optionally substituted with a substituent selected from the group consisting of halo, alkyl, alkoxy, cycloalkyl, aryl, aryloxy, arylalkyl, haloalkyl, polyhaloalkyl, haloalkoxy, polyhaloalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, arylthio, arylsulfinyl,  
30 arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, alkoxycarbonyl, aryloxycarbonyl,

heteroaryloxycarbonyl, acyl, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, carboxy, hydroxy, tetrazolyl, imidazole, oxazole, triazole, and -SOR wherein R is hydroxy, alkyl, aryl, alkoxy, aryloxy, cycloalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy or cycloheteroalkylalkoxy.

5 Preferably the linker is a methylene group, a carbonyl group, a methylene group coupled to a carbonyl group, a methylene group having a methyl substituent or a methylene group containing an optionally substituted phenyl group. The alkylene group is C1-C8, and preferably C1-C4, and more preferably C1. Preferably, haloalkyl or polyhaloalkyl groups are CF<sub>3</sub> or CF<sub>3</sub>CH<sub>2</sub> and haloalkoxy or polyhaloalkoxy groups are  
10 CF<sub>3</sub>O or CF<sub>3</sub>CH<sub>2</sub>O.

Ar<sup>1</sup> or Ar<sup>2</sup> is an aryl group which is the residue of an aromatic hydrocarbon containing one or more rings optionally including one or more heteroatoms, selected from the group of O, N and S. Preferably the aryl group has 6-12 carbon atoms and up to 3 heteroatoms, and more preferably the aryl group has 6-8 carbon atoms and 1 or 2  
15 heteroatoms. More preferably, the aryl group is a saturated or unsaturated 5-7 membered heterocycle and even more preferably a saturated or unsaturated 5-6 membered heterocycle. Most preferably, the aryl group is phenyl or residues of an optionally benzo-fused heterocycle containing up to 3 heteroatoms selected from S, N and O. The most  
20 preferable aryl groups are indolyl, isoquinolyl, quinolyl, benzimidazolyl, benzotriazolyl, benzothiazolyl, benzofuranyl, pyridyl, thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, morpholinyl or piperidyl. When the linker is not CO or an isostere thereof, Ar<sup>1</sup> or Ar<sup>2</sup> is most preferably indolyl, benzimidazolyl or benzotriazolyl.

Ar<sup>1</sup> and Ar<sup>2</sup> and/or other aryl substituents are optionally substituted with substituents including one or more of halo, alkyl, alkoxy, cycloalkyl, aryl, aryloxy,  
25 arylalkyl, haloalkyl, polyhaloalkyl, haloalkoxy, polyhaloalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, arylthio, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, acyl, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano,  
30 carboxy, hydroxy, tetrazolyl, imidazole, oxazole, triazole, and -SOR wherein R is hydroxy, alkyl, aryl, alkoxy, aryloxy, cycloalkyl, heteroaryl, heteroarylalkyl,

heteroaryloxy, heteroarylalkoxy or cycloheteroalkylalkoxy. Preferably the substituents include halo, alkoxy, alkoxyaryl, aminoalkyl, aminoaryl, and substituted aryl. Most preferably  $Ar^1$  and  $Ar^2$  include at least one halo, alkoxy, or N-containing substituent.

Y is selected from the group consisting of H, optionally substituted alkyl, alkoxy, 5 cycloalkyl, cycloheteroalkyl, aryl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, amino, and alkylamino. Preferred groups are hydroxy, alkyl and carbonyl.

The compounds of formula (2) set forth above are defined by the nature of the 10 substituents on the piperazine or piperidine ring. Piperidine rings are preferred.

In formula (2),  $Ar^1$  is optionally substituted furanyl, thiophenyl, optionally substituted phenyl systems having 0, 1, or 2 heterocyclic N atoms or naphthyl systems having 0, 1, 2 or 3 heterocyclic N atoms. The nature of the substituents and the preferred substituents are discussed below.

15  $X^1$  is CO or an isostere thereof. Thus, in addition to CO,  $X^1$  may be  $CH_2$ , SO,  $SO_2$ , or CHOH. CO is preferred.

Z is CH or N.

$X^2$  is  $CH_2$  or an isostere thereof.

The aryl moieties represented by  $Ar^1$  and Ph in formula (2) may optionally be 20 substituted by substituents including straight or branched chain alkyl (1-6C), straight or branched chain alkenyl (2-6C), halo, RCO, COOR,  $CONR_2$ ,  $SO_2NR_2$ , -OOCR, -NROCR, -NROCOR, OR, SR,  $NR_2$ ,  $NO_2$ , CN, or  $CF_3$ , wherein R is H or straight or branched chain alkyl (1-6C). Phenyl moieties may also be substituted with an additional phenyl residue, preferably at the 4-position. Any two substituents may form a 5-7 membered carbocyclic 25 or heterocyclic ring subject to the proviso. The additional phenyl residue may itself be substituted with the substituents set forth above. The additional phenyl may be substituted in all five positions, but preferably less, preferably in 1-2 positions or not at all.

Preferred substituents include halo, alkyl (1-6C), OR, SR and  $NR_2$ , more preferably halo, OR and alkyl (1-4C), most preferably halo and  $OCH_3$ . The substituents on the 30 phenyl moiety as an embodiment of  $Ar^1$  or on Ph may occupy all five available positions, preferably 1-2 positions or the phenyl is unsubstituted. If  $Ar^1$  comprises a pyridyl residue,



only 4 positions are available; preferably only 1-2 positions are substituted or preferably the pyridyl is unsubstituted. If Ar<sup>1</sup> comprises furanyl or thiophenyl, only 3 positions are available; preferred numbers of substitutions in this case are 1 or 0.

n may be 0 or 1, and is preferably 0. However, when n is 1, Y is present and may  
5 be alkyl, arylalkyl or aryl, all of which may optionally be substituted by the substituents set forth above. Preferred embodiments of Y include unsubstituted alkyl (1-6C) and unsubstituted arylalkyl (7-11C), most preferably unsubstituted lower alkyl (1-4C).

The compounds of formula (1) or (2) may be supplied in the form of their pharmaceutically acceptable acid-addition salts including salts of inorganic acids such as  
10 hydrochloric, sulfuric, hydrobromic, or phosphoric acid or salts of organic acids such as acetic, tartaric, succinic, benzoic, salicylic, and the like. If a carboxyl moiety is present on the compounds of formula (1) or (2), the compound may also be supplied as a salt with a pharmaceutically acceptable cation.

In the event the compounds of formula (1) or (2) contains one or more chiral  
15 centers, all of the stereoisomers are included within the scope of the invention, as well as mixtures thereof. Thus, the compounds of formula (1) or (2) in these instances may be supplied as a single stereoisomer, as a racemic mixture, as a partially racemic mixture, or generally, a mixture of stereoisomers in any proportion.


The invention is directed, in addition to methods of treatment, to compounds  
20 falling within the scope of formula (1) or (2) as compositions of matter.

In particular, the invention is directed to compounds of the general formula (2), and the pharmaceutically acceptable salts thereof, wherein Ar<sup>1</sup> is optionally substituted furanyl, thiophenyl, phenyl system containing 0, 1 or 2 N as heterocyclic atoms or naphthyl system containing 0, 1, 2 or 3 N as heterocyclic atoms, X<sup>1</sup> is CO or an isostere thereof; Y is  
25 optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C); n is 1; Z is N or CH; X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof; and Ph is optionally substituted phenyl.

The invention is also directed to compounds of formula (2) and the pharmaceutically acceptable salts thereof, wherein Ar<sup>1</sup> is optionally substituted furanyl,  
30 thiophenyl, phenyl system containing 1 or 2 N atoms as heterocyclic atoms or naphthyl system containing 0, 1, 2, or 3 N atoms as heterocyclic atoms, X<sup>1</sup> is CO or an isostere

thereof; Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C); n is 0-1; Z is N or CH; X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof; and Ph is optionally substituted phenyl; wherein said substituents on Ar<sup>1</sup> and Ph are independently selected from the group consisting of straight or branched chain alkyl (1-6C), straight or branched chain alkenyl (2-6C), halo, RCO, COOR, CONR<sub>2</sub>, SO<sub>2</sub>NR<sub>2</sub>, -OOCR, -NROCR, -NROCOR, OR, SR, NR<sub>2</sub>, NO<sub>2</sub>, CN, or CF<sub>3</sub>, wherein R is H or straight or branched chain alkyl (1-6C).

The invention is also directed to a compound of the formula (2) and the pharmaceutically acceptable salts thereof, wherein Ar<sup>1</sup> is an ortho substituted furanyl, thiophenyl, phenyl system containing 0, 1 or 2 N as heterocyclic atoms, or naphthyl system containing 0, 1, 2 or 3 N as heterocyclic atoms; said ortho substituent is straight or branched chain alkyl (1-6C), straight or branched chain alkenyl (2-6C), halo, RCO, COOR, CONR<sub>2</sub>, SO<sub>2</sub>NR<sub>2</sub>, -OOCR, -NROCR, -NROCOR, OR, SR, NR<sub>2</sub>, NO<sub>2</sub>, CN, or CF<sub>3</sub>, wherein R is H or straight or branched chain alkyl (1-6C), with the proviso that when said Ar<sup>1</sup> is phenyl and said ortho substituent is OR, either R must be alkyl (3-6C) or Ar<sup>1</sup> must be



where R' is H, alkyl (1-6C), alkenyl (2-6C) or arylalkyl (7-12C);

X<sup>1</sup> is CO or an isostere thereof; Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C); n is 0-1; Z is N or CH; and Ph is optionally substituted phenyl.

The ortho position is defined herein as the position in the ring adjacent the X<sup>1</sup> position, for example the 2-position in a 5-membered ring.

Preferred embodiments of the compounds of the invention are as described above with respect to compounds useful in the invention methods. In particular piperidine forms are preferred over piperazines.

Particularly preferred compounds of the invention include:





1-(2-methoxy-4-hydroxybenzoyl)-4-benzylpiperidine;






1-(2-methoxy-4-methoxybenzoyl)-4-benzylpiperidine;








1-(2-methoxy-4-benzyloxybenzoyl)-4-benzylpiperidine; and

1-(2-methoxy-4-methoxybenzoyl)-4-(4-fluorobenzyl)piperidine.


In addition, the following table illustrates preferred compounds of the invention.

Compound #	Structure
1	
2	
3	
4	


5	 <chem>O=C1CCCC1N2CCN(C(=O)c3ccc4[nH]cnc4c3)CC2</chem>
6	 <chem>O=C1CCCCC1N2CCN(C(=O)c3ccc4[nH]cnc4c3)CC2</chem>
7	 <chem>O=C1CCCC1CC(=O)N2CCN(C(=O)c3ccc4[nH]cnc4c3)CC2</chem>
8	 <chem>CCCC(C)C(=O)N2CCN(C(=O)c3ccc4[nH]cnc4c3)CC2</chem>
9	 <chem>O=C1CCCC1CC(=O)N2CCN(C(=O)c3ccc4[nH]cnc4c3)CC2</chem>




17




18




19




20




21




22




23




24










25














26




27	
28	
29	
30	
31	
32	
33	




34	 <chem>COc1ccc(cc1)C(=O)N2C=CC3=CC=C(C=C3C2=CC=C3)C(=O)N4CCc5ccccc5CC4</chem>
35	 <chem>COc1ccc(cc1)C(=O)N2C=CC3=CC=C(C=C3C2=CC=C3)C(=O)N4CCc5ccccc5CC4</chem>
36	 <chem>COc1ccc(cc1)C(=O)N2C=CC3=CC=C(C=C3C2=CC=C3)C(=O)N4CCc5ccccc5CC4</chem>
37	 <chem>COc1ccc(cc1)C(=O)N2C=CC3=CC=C(C=C3C2=CC=C3)C(=O)N4CCc5ccccc5CC4</chem>
38	 <chem>COc1ccc(cc1)C(=O)N2C=CC3=CC=C(C=C3C2=CC=C3)C(=O)N4CCc5ccccc5CC4</chem>
39	 <chem>COc1ccc(cc1)C(=O)N2C=CC3=CC=C(C=C3C2=CC=C3)C(=O)N4CCc5ccccc5CC4</chem>

40	 <chem>O=C(Cc1ccccc1)N2CCN(C(=O)c3ccc4ncnc4c3)CC2</chem>
41	 <chem>CC(=O)OC(=O)N2CCN(C(=O)c3ccc4ncnc4c3)CC2</chem>
42	 <chem>O=C(OCC1=CC=CC=C1)N2CCN(C(=O)c3ccc4ncnc4c3)CC2</chem>
43	 <chem>O=C(Cc1ccc(OC)cc1)N2CCN(C(=O)c3ccc4ncnc4c3)CC2</chem>
44	 <chem>CCOC(=O)C(c1ccccc1)C(=O)N2CCN(C(=O)c3ccc4ncnc4c3)CC2</chem>


45




46




47


















48











49




50	
51	
52	
53	
54	
55	
56	


57	
58	
59	
60	
61	
62	
63	
64	

65	
66	
67	
68	
69	
70	
71	
72	


73




74




75




76




77




78




79




80




81




82




83




84



85




86







87




88




89




90




91




92




93




94




95




96




97




98




99




100




101




102










103



104



105	
106	
107	
108	
109	
110	




The invention is also directed to methods to prepare the compounds useful in the invention by forming the carboxamides from the appropriate aroyl compounds as halides  
5 or free acids.


### Synthesis Methods

Preferably, the compounds useful in the invention are synthesized, generally, by coupling an aroyl moiety to a benzyl-substituted piperazine or piperidine. The general  
10 approach is shown in Reaction Schemes 1-6 hereinbelow. Reaction Scheme 3 is directed to the particular circumstance wherein Ar<sup>2</sup> includes an amino group to be functionalized in the presence of potentially competing groups.


#### **Reaction Scheme 1**



## Reaction Scheme 2




## Reaction Scheme 3




5


## Reaction Scheme 4



## Reaction Scheme 5



## Reaction Scheme 6



5

Administration and Use

The methods of the invention are directed to treating conditions associated with p38 kinase activity, for example, inflammatory conditions. Thus, the compounds of formula (1) or their pharmaceutically acceptable salts are used in the manufacture of a medicament for prophylactic or therapeutic treatment of mammals, including humans, in respect of conditions characterized by excessive production of cytokines and/or

inappropriate or unregulated cytokine activity on such cells as cardiomyocytes, cardiofibroblasts and macrophages.

The compounds of formulas (1) and (2) inhibit the production of cytokines such as TNF, IL-1, IL-6 and IL-8, cytokines that are important proinflammatory constituents in many different disease states and syndromes. Thus, inhibition of these cytokines has benefit in controlling and mitigating many diseases. The compounds of formulas (1) and (2) are shown herein to inhibit a member of the MAP kinase family variously called p38 MAPK (or p38), CSBP, or SAPK-2. The activation of this protein has been shown to accompany exacerbation of the diseases in response to stress caused, for example, by treatment with lipopolysaccharides or cytokines such as TNF and IL-1. Inhibition of p38 activity, therefore, is predictive of the ability of a medicament to provide a beneficial effect in treating diseases such as coronary artery disease, congestive heart failure, cardiomyopathy, myocarditis, vasculitis, restenosis, such as occurs following coronary angioplasty, atherosclerosis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, multiple sclerosis, acute respiratory distress syndrome (ARDS), asthma, chronic obstructive pulmonary disease (COPD), silicosis, pulmonary sarcosis, sepsis, septic shock, endotoxic shock, toxic shock syndrome, heart and brain failure (stroke) that are characterized by ischemia and reperfusion injury, surgical procedures, such as transplantation procedures and graft rejections, cardiopulmonary bypass, coronary artery bypass graft, CNS injuries, including open and closed head trauma, inflammatory eye conditions such as conjunctivitis and uveitis, acute renal failure, glomerulonephritis, inflammatory bowel diseases, such as Crohn's disease or ulcerative colitis, graft vs host disease, bone resorption diseases like osteoporosis, type II diabetes, pyresis, psoriasis, cachexia, viral diseases such as those caused by HIV, CMV, and Herpes, and cerebral malaria.

Within the last several years, p38 has been shown to comprise a group of MAP kinases designated p38 $\alpha$ , p38 $\beta$ , p38 $\gamma$  and p38 $\delta$ . Jiang, Y. *et al. J Biol Chem* (1996) 271:17920-17926 first reported characterization of p38 $\beta$  as a 372-amino acid protein closely related to p38 $\alpha$ . Kumar, S. *et al. Biochem Biophys Res Comm* (1997) 235:533-538 and Stein, B. *et al. J Biol Chem* (1997) 272:19509-19517 reported a second isoform of p38 $\beta$ , p38 $\beta$ 2 containing 364 amino acids with 73% identity to p38 $\alpha$ . All of these reports



show evidence that p38 $\beta$  is activated by proinflammatory cytokines and environmental stress, although the second reported p38 $\beta$  isoform, p38 $\beta$ 2, appears to be preferentially expressed in the CNS, heart and skeletal muscle compared to the more ubiquitous tissue expression of p38 $\alpha$ . Furthermore, activated transcription factor-2 (ATF-2) was observed  
5 to be a better substrate for p38 $\beta$ 2 than for p38 $\alpha$ , thus suggesting that separate mechanisms of action may be associated with these forms. The physiological role of p38 $\beta$ 1 has been called into question by the latter two reports since it cannot be found in human tissue and does not exhibit appreciable kinase activity with the substrates of p38 $\alpha$ .

The identification of p38 $\gamma$  was reported by Li, Z. *et al. Biochem Biophys Res*  
10 *Comm* (1996) 228:334-340 and of p38 $\delta$  by Wang, X., *et al., J Biol Chem* (1997) 272:23668-23674 and by Kumar, S., *et al., Biochem Biophys Res Comm* (1997) 235:533-538. The data suggest that these two p38 isoforms ( $\gamma$  and  $\delta$ ) represent a unique subset of the MAPK family based on their tissue expression patterns, substrate utilization, response to direct and indirect stimuli, and susceptibility to kinase inhibitors.

15 Various results with regard to response to drugs targeting the p38 family as between p38 $\alpha$  and either the putative p38 $\beta$ 1 or p38 $\beta$ 2 or both were reported by Jiang, Kumar, and Stein cited above as well as by Eyers, P.A. *et al. Chem and Biol* (1995) 5:321-328. An additional paper by Wang, Y. *et al. J Biol Chem* (1998) 273:2161-2168 suggests the significance of such differential effects. As pointed out by Wang, a number of stimuli,  
20 such as myocardial infarction, hypertension, valvular diseases, viral myocarditis, and dilated cardiomyopathy lead to an increase in cardiac workload and elevated mechanical stress on cardiomyocytes. These are said to lead to an adaptive hypertrophic response which, if not controlled, has decidedly negative consequences. Wang cites previous studies which have shown that in ischemia reperfusion treated hearts, p38 MAPK activities  
25 are elevated in association with hypertrophy and programmed cell death. Wang shows in the cited paper that activation of p38 $\beta$  activity results in hypertrophy, whereas activation of p38 $\alpha$  activity leads to myocyte apoptosis. Thus, selective inhibition of p38 $\alpha$  activity as compared to p38 $\beta$  activity will be of benefit in treating conditions associated with cardiac failure. These conditions include congestive heart failure, cardiomyopathy, myocarditis,  
30 vasculitis, vascular restenosis, valvular disease, conditions associated with cardiopulmonary bypass, coronary artery bypass, grafts and vascular grafts. Further, to the

extent that the  $\alpha$ -isoform is toxic in other muscle cell types,  $\alpha$ -selective inhibitors would be useful for conditions associated with cachexia attributed to TNF or other conditions such as cancer, infection, or autoimmune disease.

Thus, the invention is directed to the use of the compounds of formulas (1) and (2) which selectively inhibit the activity of the p38 $\alpha$  isoform for treating conditions associated with activation of p38 $\alpha$ , in particular those associated with cardiac hypertrophy, ischemia or other environmental stress such as oxidation injury, hyperosmolarity or other agents or factors that activate p38 $\alpha$  kinase, or cardiac failure, for example, congestive heart failure, cardiomyopathy and myocarditis.

The manner of administration and formulation of the compounds useful in the invention will depend on the nature of the condition, the severity of the condition, the particular subject to be treated, and the judgement of the practitioner; formulation will depend on mode of administration. As the compounds of formulas (1) and (2) are small molecules, they are conveniently administered by oral administration by compounding them with suitable pharmaceutical excipients so as to provide tablets, capsules, syrups, and the like. Suitable formulations for oral administration may also include minor components such as buffers, flavoring agents and the like. Typically, the amount of active ingredient in the formulations will be in the range of 5%-95% of the total formulation, but wide variation is permitted depending on the carrier. Suitable carriers include sucrose, pectin, magnesium stearate, lactose, peanut oil, olive oil, water, and the like.

The compounds of formulas (1) and (2) may also be administered through suppositories or other transmucosal vehicles. Typically, such formulations will include excipients that facilitate the passage of the compound through the mucosa such as pharmaceutically acceptable detergents.

The compounds may also be administered topically, for topical conditions such as psoriasis, or in formulation intended to penetrate the skin. These include lotions, creams, ointments and the like which can be formulated by known methods.

The compounds may also be administered by injection, including intravenous, intramuscular, subcutaneous or intraperitoneal injection. Typical formulations for such use are liquid formulations in isotonic vehicles such as Hank's solution or Ringer's solution.

Alternative formulations include nasal sprays, liposomal formulations, slow-release formulations, and the like, as are known in the art.

Any suitable formulation may be used. A compendium of art-known formulations is found in Remington's Pharmaceutical Sciences, latest edition, Mack Publishing

5 Company, Easton, PA. Reference to this manual is routine in the art.

The dosages of the compounds of formula (1) will depend on a number of factors which will vary from patient to patient. However, it is believed that generally, the daily oral dosage will utilize 0.001-100 mg/kg total body weight, preferably from 0.01-50 mg/kg and more preferably about 0.01 mg/kg-10 mg/kg. The dose regimen will vary,  
10 however, depending on the conditions being treated and the judgment of the practitioner.

The inhibitors of p38 kinase can be used as single therapeutic agents or in combination with other therapeutic agents. Drugs that could be usefully combined with these compounds include natural or synthetic corticosteroids, particularly prednisone and its derivatives, monoclonal antibodies targeting cells of the immune system, antibodies or  
15 soluble receptors or receptor fusion proteins targeting immune or non-immune cytokines, and small molecule inhibitors of cell division, protein synthesis, or mRNA transcription or translation, or inhibitors of immune cell differentiation or activation.

As implicated above, although the compounds of formula (1) may be used in humans, they are also available for veterinary use in treating animal subjects.

20


The following examples are intended to illustrate but not to limit the invention.

#### Example 1

##### Preparation of 4-benzylpiperidin-1-yl benzene carboxamide

25

##### (1-benzoyl-4-benzyl piperidine)



This example illustrates Reaction Scheme 1.


To a mixture of 4-benzylpiperidine (0.263 g, 1.5 mM) and diisopropylethyl-amine (0.53 mL, 3.0 mM) in 10 mL dichloromethane at room temperature was added benzoyl chloride (0.281 g, 2.0 mM) and the reaction mixture was stirred at room temperature for 20 h. The reaction mixture was poured into water and extracted with ethyl acetate (2 x 25 mL). The combined organic extract was washed with 1 N HCl, water and brine. The extract was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The residue was chromatographed on a column of silica gel with ethyl acetate-hexane (10 to 50%, gradient). Evaporation of the desired fractions gave 0.402 g (96%) of the title compound as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.10-7.40 (m, 10H), 4.60-4.70 (m, 1H), 3.70-3.80 (m, 1H), 2.80-2.90 (m, 1H), 2.70-2.80 (m, 1H), 2.60 (d, 2H), 1.70-1.80 (m, 1H), 1.50-1.70 (m, 2H), 1.20-1.40 (m, 2H). MS (ESI) *m/e* 279 (M<sup>+</sup>)

### Example 2

#### Preparation of 4-Benzylpiperidin-1-yl-(4-amino)benzene carboxamide

#### (1-(4-aminobenzoyl)-4-benzyl piperidine)

This example illustrates Reaction Scheme 2:



4-Aminobenzoic acid (1.61g, 10 mM) was reacted with EDAC (1.92g, 10 mM) in 40 mL dry DMF for 15 minutes. 4-Benzylpiperidine (1.75g, 10 mM) was added followed by DMAP (20 mg, catalyst) and the reaction mixture was stirred for 20 h. The mixture was poured into water and extracted with methylene chloride (3x100 mL). The combined extract was washed with dilute hydrochloric acid, saturated sodium bicarbonate and water and dried over MgSO<sub>4</sub>. After evaporation of the solvent, the residue was chromatographed with methylene chloride-methanol (0 to 2% methanol, gradient) to yield 1.60 g (50%) of the title compound after recrystallization from ether-Hexane. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.40-7.10 (m, 7 H), 6.70 (d, 2 H), 3.95-3.90 (broad m, 2 H), 2.85-2.75 (broad m, 2 H), 2.6 (d, 2

H), 1.85-1.80 (m, 1 H), 2.75-2.65 (broad m, 2 H), 1.15-1.05 (broad m, 2 H). MS (ESI) m/e 294 (M<sup>+</sup>).

Examples 3 and 4 illustrate Reaction Scheme 3:

5

### Example 3

#### Preparation of 1-(2-chloro-4-aminobenzoyl)-4-benzyl piperidine

##### A. Preparation of the BOC-protected Intermediate

To a solution of 2-Chloro-4-*t*-butoxycarbonylaminobenzoic acid (0.50 g, 1.85 mM) in CH<sub>2</sub>Cl<sub>2</sub> (8.0 mL) was added triethylamine (0.37 g, 0.52 mL, 3.70 mM) and 4-benzylpiperidine (0.65 g, 0.65 mL, 3.70 mM). This was followed by the dropwise addition of thionyl chloride (0.44 g, 0.27 mL, 3.70 mM). The mixture was allowed to stir at room temperature for 1 h, whereupon it was poured into water and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 10 mL). The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. Silica gel chromatography (5% methanol in dichloromethane) afforded 0.30 g (38%) of BOC-protected intermediate, i.e., 1-(2-chloro-4-*t*-butoxycarbonyl aminobenzoyl)-4-benzyl piperidine. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.30-7.10 (m, 2 H), 6.60 (s, 1 H), 4.70 (m, 1 H), 3.30 (m, 1 H), 2.50 (d, 2 H), 1.80 (m, 2 H), 1.60 (m, 2 H), 1.50 (s, 9 H), 1.20 (m, 3 H). MS (ESI) m/e 428 (M<sup>+</sup>).

B. 1-(2-chloro-4-*t*-butoxycarbonyl aminobenzoyl)-4-benzyl piperidine (0.15 g, 0.35 mM) from paragraph A was added to 4.0 N HCl in dioxane (2.0 mL). The mixture was stirred for 2 hours at room temperature whereupon it was triturated with Et<sub>2</sub>O. The Et<sub>2</sub>O was decanted and the residue was dried in vacuo to yield 0.10 g (87%) of the title compound. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.50 (m, 3 H), 7.30 (m, 4 H), 7.20 (d, 1 H), 7.10 (d, 2 H), 4.6 (t, 1 H), 3.30 (d, 1 H), 2.90 (t, 1 H), 2.70 (t, 1 H), 2.50 (s, 2 H), 1.70 (d, 2 H), 1.60 (t, 1 H), 1.20 (broad s, 2 H). MS (ESI) m/e 327 (M<sup>+</sup>-1).

### Example 4

#### Preparation of 1-(2-chloro-4-methylaminobenzoyl)-4-benzyl piperidine

A solution of 1-(2-chloro-4-*t*-butoxycarbonyl aminobenzoyl)-4-benzyl piperidine (0.88g, 2.06 mM) in THF (4.0 mL) from paragraph A of Example 3 was added dropwise to an ice-cooled suspension of NaH in THF (4.0 mL). Stirring was continued for 0.5 hours

before adding iodomethane (0.35 g, 0.15 mL, 2.47 mM) dropwise. The mixture was then allowed to stir at room temperature for 16 hours, at which time it was poured into water and extracted with ethyl acetate (3 x 10 mL). The combined extracts were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was dissolved in 4.0 N HCl in dioxane (10.0 mL) and stirred at room temperature for 2 hours whereupon it was triturated with Et<sub>2</sub>O. The Et<sub>2</sub>O was decanted and the residue was dried in vacuo to yield 0.56 g (79%) of 1-(2-chloro-4-methylaminobenzoyl)-4-benzyl piperidine. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.60-7.10 (m, 8 H), 4.70 (t, 1 H), 3.40 (broad s, 1 H), 3.00 (m, 1 H), 2.90 (s, 3 H), 2.70 (m, 1 H) 2.60 (m, 1 H), 1.80 (broad s, 2 H), 1.60 (broad s, 1 H), 1.30 (m, 2 H). MS (ESI) m/e 342 (M<sup>+</sup>).

10

### Example 5

#### Preparation of Additional Compounds of Formula (2)

Using the techniques set forth in Examples 1-4, the following additional compounds of the invention were prepared:

- 15 1-(4-methoxybenzoyl)-4-benzyl piperidine,
- 1-(4-cyanobenzoyl)-4-benzyl piperidine,
- 1-(4-phenylbenzoyl)-4-benzyl piperidine,
- 1-(3-methoxybenzoyl)-4-benzyl piperidine,
- 1-(3,5-dimethoxybenzoyl)-4-benzyl piperidine,
- 20 1-(2-chlorobenzoyl)-4-benzyl piperidine,
- 1-(2-bromobenzoyl)-4-benzyl piperidine,
- 1-(2-iodobenzoyl)-4-benzyl piperidine,
- 1-(2-methoxybenzoyl)-4-benzyl piperidine,
- 1-(2-methylbenzoyl)-4-benzyl piperidine,
- 25 1-(2-phenylbenzoyl)-4-benzyl piperidine,
- 1-(2-(2-carbomethoxyethen-1-yl)benzoyl)-4-benzyl piperidine,
- 1-(2-naphthoyl)-4-benzyl piperidine,
- 1-(1-naphthoyl)-4-benzyl piperidine,
- 1-(2-furanoyl)-4-benzyl piperidine,
- 30 1-(2-thiophenoyl)-4-benzyl piperidine,
- 1-(2-dimethylaminobenzoyl)-4-benzyl piperidine,

1-(3-dimethylaminobenzoyl)-4-benzyl piperidine,  
1-(4-dimethylaminobenzoyl)-4-benzyl piperidine,  
1-(3,4,5-trimethoxybenzoyl)-4-benzyl piperidine,  
1-(2-acetamidobenzoyl)-4-benzyl piperidine,  
5 1-(3-acetamidobenzoyl)-4-benzyl piperidine,  
1-(4-acetamidobenzoyl)-4-benzyl piperidine, and  
1-(4-formamidobenzoyl)-4-benzyl piperidine.

### Example 6

#### 10 Preparation of 1-Phenyl-8-1,3,8-triazaspiro[4.5]decan-4-one-3-acetyl-4-benzylpiperidine

This example illustrates Reaction Scheme 4.

Acid I [*J Med Chem* (1996), 39(16):3169-3173]: 1.0 g, 2.57 mmol and 4-Benzylpiperidine, (0.46 g, 2.6 mmol) were treated with 0.51 g EDAC.HCl in 10 mL dichloromethane (DCM) in the presence of 200  $\mu$ L 0.1M DMAP in DMF. After stirring  
15 at room temperature for 4 hours, the reaction mixture was concentrated and the residue was taken up in 50 mL ethyl acetate (EtOAc). The EtOAc layer was washed with 10% aqueous sodium carbonate, 10% aqueous hydrochloric acid, sodium chloride solution and dried over anhydrous sodium sulfate. Concentration gave crude product as a white solid. This material was chromatographed using ethylacetate/hexane on silica gel to the give  
20 Boc-protected compound as a white solid. 700 mg EIMS  $M^+$  546. 100 mg of the Boc-protected compound was treated with 10 mL 4N HCl in dioxane for 60 mins. The reaction mixture was concentrated and the solid obtained was filtered through a sintered glass funnel and washed with hexane. Extensive drying gave the hydrochloride salt of Example 1. EIMS  $M^+$  446. 65 mg.

25

### Example 7

#### Preparation of 4-Phenyl-3-pyrrolyl-4-benzylpiperidine

This example illustrates Reaction Scheme 4.

4-Phenyl-3-pyrrole carboxylic acid (333 mg, 3mole) and 4-Benzylpiperidine  
30 (525mg 3mmole) were dissolved in 15ml dichloromethane/dimethylformamide(1.1). EDAC-HCl ( 573mg, 3mmole) was added and the reaction mixture was stirred overnight.

The solvent was removed and residue redissolved in EtOAc, the EtOAc was washed with 10% HCl, 0.1M NaOH, water, the EtOAc extract was dried over magnesium sulfate (anh). Evaporated EtOAc to obtain 489mg product.

Unless otherwise indicated all compounds were prepared following the method outlined above using carboxylic acids that are either commercially available or are available by synthesis, using methods known in the literature.

### Example 8

#### Preparation of (V) 4-(4')-Nitrophenyl-3-purroyl-4-benzylpiperidine

10 This example illustrates Reaction Scheme 6.

Preparation of II: Sodium hydride (614mg 25.6mmole) was suspended in 40ml diethyl ether. Methyl 4-nitrophenyl-trans-cinnamate (4.14g, 20mmole), tosylmethylisocyanate (4.29g, 22mole) dissolved in Et<sub>2</sub>O/DMSO (2:1) 40ml were added dropwise under nitrogen over 20min period. The mixture was stirred for 1 hour at room temperature. Water was added to the reaction mixture and precipitate was filtered. The aqueous layer was extracted with Et<sub>2</sub>O (2 x 75ml), The combined extracts were dried over MgSO<sub>4</sub> and solvent removed under reduced pressure to give 1.42g product. EIMS M<sup>+</sup> 247.

Preparation of III: Methyl-3-(4-nitrophenyl)-2-pyrrole carboxylate (626mg 2.54mmole) was treated with potassium hydroxide (713mg, 12.7mmole) in methanol/water (50:50). The reaction mixture was refluxed 2.5hours. Methanol was evaporated, the solution was cooled and acidified with HCl(aq) to pH 7. The mixture was extracted with ethyl acetate, combined extracts dried over MgSO<sub>4</sub>(anh.) and concentrated to give 440mg product. EIMS M<sup>+</sup> 232.

25 Preparation of IV: 3-(4-nitrophenyl)-2-pyrrole carboxylic acid (440mg 1.89mmole) and 4-Benzylpiperidine (331mg 1.89mmole) were dissolved in DCM/DMF(10:1), EDAC (360mg, 1.89mmole) was added and the reaction mixture was stirred for 18hr at room temperature. The solvent was evaporated, residue dissolved in EtOAc, washed with 10% HCl, 0.10M NaOH, water, the EtOAc layer was dried over MgSO<sub>4</sub>, solvent evaporated to give 240mg product. Product was chromatographed on silica gel with EtOAc:hex (8:2) to give 180mg purified product. EIMS M<sup>+</sup> 389.



Preparation of (V) 4-(4')-Nitrophenyl-3-purroyl-4-benzylpiperidine: 180mg of the nitro compound was dissolved in 10ml methanol, 10% Pd/C (60mg) was added and hydrogenated under hydrogen balloon for 2 hours. The catalyst was filtered over celite, solvent was evaporated to give 136mg of the desired product. EIMS M<sup>+</sup> 359.

5

### Example 9

#### Assay for p38 Kinase Inhibition

The compounds to be tested were solubilized in DMSO and diluted into water to the desired concentrations. The p38 kinase was diluted to 10 µg/ml into a buffer  
10 containing 20 mM MOPS, pH 7.0, 25 mM beta-glycerol phosphate, 2 mg/ml gelatin, 0.5 mM EGTA, and 4 mM DTT.

The reaction was carried out by mixing 20 µl test compound with 10 µl of a substrate cocktail containing 500 µg/ml peptide substrate and 0.2 mM ATP (+200 µCi/ml gamma-32P-ATP) in a 4x assay buffer. The reaction was initiated by the addition of 10 µl  
15 of p38 kinase. Final assay conditions were 25 mM MOPS, pH 7.0, 26.25 mM beta-glycerol phosphate, 80 mM KCl, 22 mM MgCl<sub>2</sub>, 3 mM MgSO<sub>4</sub>, 1 mg/ml gelatin, 0.625 mM EGTA, 1 mM DTT, 125 µg/ml peptide substrate, 50 µM ATP, and 2.5 µg/ml enzyme. After a 40 minute incubation at room temperature, the reaction was stopped by the addition of 10 µl per reaction of 0.25 M phosphoric acid.

20 A portion of the reaction was spotted onto a disk of P81 phosphocellulose paper, the filters were dried for 2 minutes and then washed 4x in 75 mM H<sub>3</sub>PO<sub>4</sub>. The filters were rinsed briefly in 95% ethanol, dried, then placed in scintillation vials with liquid scintillation cocktail.

Alternatively, the substrate is previously biotinylated and the resulting reactions  
25 are spotted on SAM<sup>2</sup>™ streptavidin filter squares (Promega). The filters are washed 4x in 2M NaCl, 4x in 2M NaCl with 1% phosphoric acid, 2x in water, and briefly in 95% ethanol. The filter squares are dried and placed in scintillation vials with liquid scintillation cocktail.

Counts incorporated are determined on a scintillation counter. Relative enzyme  
30 activity is calculated by subtracting background counts (counts measured in the absence of enzyme) from each result, and comparing the resulting counts to those obtained in the

absence of inhibitor.  $IC_{50}$  values were determined with curve-fitting plots available with common software packages. Approximate  $IC_{50}$  values were calculated using formula

$$IC_{50} (app) = A \times i / (1 - A)$$


where A=fractional activity and i=total inhibitor concentration.

- 5        The percent inhibition of p38- $\alpha$  kinase was determined at various concentrations in order to determine  $IC_{50}$  values. Table 1 shows compounds tested for the percent inhibition of p38- $\alpha$  kinase at concentrations of 15  $\mu$ M, 5  $\mu$ M, 1  $\mu$ M and 0.2  $\mu$ M. The table shows various embodiments of  $Ar^1X^1$ ; unless otherwise noted, Ph is unsubstituted phenyl, Z is CH and n is 0. Thus, all of the compounds in Table 1 are 4-benzyl piperidines, unless
- 10       otherwise noted. All show substantial inhibition at 15  $\mu$ M, some as high as 99%. Virtually all are inhibitory at 0.2  $\mu$ M.


Table 1	
Comp. No.	$Ar^1X^1$
1	benzoyl
2	4-methoxybenzoyl
3	4-cyanobenzoyl
4	4-phenylbenzoyl
5	3-methoxybenzoyl
6	3,5-dimethoxybenzoyl
7	2-chlorobenzoyl
8	2-bromobenzoyl
9	2-iodobenzoyl
10	2-methoxybenzoyl
11	2-methylbenzoyl
12	2-(2-carbomethoxy ethen-1-yl) benzoyl
13	2-naphthoyl
14	1-naphthoyl
15	2-furanoyl
16	2-thiophenoyl
17	note 1
18	2-dimethylaminobenzoyl
19	3-dimethylaminobenzoyl
20	4-dimethylaminobenzoyl
21	2-pyridoyl
22	3-pyridoyl

Table 1	
Comp. No.	Ar <sup>1</sup> X <sup>1</sup>
23	4-pyridoyl
24	note 2
25	3,4,5-trimethoxybenzoyl
26	2-chloro-4-acetamidobenzoyl
27	2-4-dimethylbenzoyl
28	4-acetamidobenzoyl
29	4-formamidobenzoyl
30	2-aminobenzoyl
31	3-aminobenzoyl
32	4-aminobenzoyl
33	4,5-dimethoxybenzoyl
34	2,4-dimethoxybenzoyl
35	2-hydroxybenzoyl
36	2-benzoyloxybenzoyl
37	3,5-bis-trifluoromethylbenzoyl
38	2,6-dimethoxybenzoyl
39	2-methylaminobenzoyl
40	6-quinolyl
41	2-chloro-4-benzamidoyl
42	2-hydroxy-4-aminobenzoyl
43	2-chloro-4-aminobenzoyl
44	2-chloro-4-methylaminobenzoyl
45	2-chloro-4-dimethylaminobenzoyl
46	2-methoxy-4-nitrobenzoyl
47	2-methoxy-4-aminobenzoyl
48	4-guanidinobenzoyl
49	2-bromobenzoyl (note 3)
50	phenylsulfonyl
51	4-amino-3-pyridoyl
52	2-methoxybenzoyl (note 4)

Note 1: In compound 17, X<sup>1</sup> is CO and Ar<sup>1</sup> has the structure:



Note 2: In compound 24, X<sup>1</sup> is CO and Ar<sup>1</sup> has the structure:



Note 3: In compound 49, n is 1 and Y is methyl. The methyl and benzyl group at the 4-position of the piperidine ring are *cis* to each other.

Note 4: In compound 52, Z is N; compound 52 is a piperazine derivative.

In some cases,  $IC_{50}$  values have been determined as well as the ratio of  $IC_{50}$  values for inhibition of  $\beta$  as compared to  $\alpha$  p38 kinase. The compounds in Table 2 show  $IC_{50}$  of about 200 nM to 1.5  $\mu$ M. Those tested are specific for p38- $\alpha$  at least by a factor of 5.

Table 2	
Comp. No.	Ar <sup>1</sup> X <sup>1</sup>
1	benzoyl
2	4-methoxybenzoyl
5	3-methoxybenzoyl
7	2-chlorobenzoyl
8	2-bromobenzoyl
9	2-iodobenzoyl
10	2-methoxybenzoyl
11	2-methylbenzoyl
16	2-thiophenoyl
18	2-dimethylaminobenzoyl
19	3-dimethylaminobenzoyl
20	4-dimethylaminobenzoyl
28	4-acetamidobenzoyl
29	4-formamidobenzoyl
32	4-aminobenzoyl
34	2,4-dimethoxybenzoyl
35	2-hydroxybenzoyl
38	2,6-dimethoxybenzoyl
39	2-methylaminobenzoyl
43	2-chloro-4-aminobenzoyl
44	2-chloro-4-methylaminobenzoyl
45	2-chloro-4-dimethylaminobenzoyl
49	2-bromobenzoyl (note 3)


### Example 10

#### Preparation of Ortho-Substituted Aryl Derivatives

The aroyl piperidines or piperazines of formula (2) can be provided with ortho-


- 5 Bindal, R.D. *et al.*, *J Org Chem* (1987) 52:3181-3185; Gshwend, H. *et al.*, *Org Rxns* (1979) 26:1. The general procedure is shown as follows:

#### Reaction Scheme 7



- 10 The starting material is first metallated with butyl lithium in the presence of base and then treated with an electrophile to provide the appropriate substitution. The electrophile may be an alkyl group, but additional functionality may be introduced by appropriate choice of the electrophile. For example, use of benzaldehyde as the source of the electrophile results in the secondary alcohol.

- 15 Compounds with further substitutions on the aroyl moiety can also be used as substrates, but those that contain an N-H require two equivalents of the butyl lithium/base.

Example 11APreparation of 4-benzylpiperidin-1-yl-(2-methoxy-4-benzyloxy)benzene carboxamide

A. Methyl 2-hydroxy-4-benzyloxybenzoate: To a solution of methyl 2,4-dihydroxybenzoate (17.5 g, 104 mM) in acetone at 0 °C were added potassium carbonate (52.5 g, 350 mM) and benzyl bromide (13.8 mL, 116 mM). The mixture was stirred at 0 °C for 2 h, followed by stirring at RT for 16 h. This mixture was filtered through a pad of Celite and concentrated. The resulting material was partitioned between ethyl acetate and water. The organic layer was washed with a saturated sodium bicarbonate solution, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The residue was purified via silica gel chromatography with ethyl acetate-hexane (1 to 20 %, gradient). Evaporation of the desired fractions resulted in 13.48 g (50 %) of the title compound as a white solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 7.80 (d, 1 H), 7.50-7.30 (m, 5 H), 6.55 (m, 2 H), 5.10 (s, 2 H), 3.95 (s, 3 H). MS (ESI) *m/e* 258 (m<sup>+</sup>).


B. Methyl 2-methoxy-4-benzyloxybenzoate: To an ice-cooled solution of methyl 2-hydroxy-4-benzyloxybenzoate from paragraph A (12.92 g, 50.1 mM) in DMF (200 mL) was added NaH in a 60 % oil dispersion (3.00 g, 75.0 mM). The mixture was stirred for 5 min, at which time MeI (7.9 mL, 127 mM) was added dropwise. The mixture was allowed to stir at RT for 16 h, whereupon it was poured onto ice and extracted with ethyl acetate. The organics were washed with water and brine, then dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to yield 13.11 g (96.3 %) of the title compound as a tan solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 7.85 (d, 1 H), 7.50-7.30 (m, 5 H), 6.55 (m, 2 H), 5.15 (s, 2 H), 3.95 (s, 3 H), 3.90 (s, 3 H). MS (ESI) *m/e* 272 (m<sup>+</sup>).

C. Title Compound: To a solution of 4-benzylpiperidine (27.5 mM, 4.85 mL) in toluene (75 mL) at -78 °C was added Me<sub>3</sub>Al (2 M solution in hexane, 32.5 mM, 16.25 mL). The mixture was stirred at -78 °C for 10 min followed by stirring at RT for 1 h.

The mixture was once again cooled to  $-78^{\circ}\text{C}$  and a solution of methyl 2-methoxy-4-benzyloxybenzoate from paragraph B (20.0 mM, 6.96 g) in toluene (25 mL) was added dropwise. Stirring was continued at  $-78^{\circ}\text{C}$  for 10 min, at which time the mixture was allowed to warm to RT and then refluxed for 16 h. The mixture was then cooled and  
5 quenched with water, made basic (pH 12) with 1 M NaOH, and stirred at RT for 0.5 h. The mixture was then extracted with ethyl acetate, washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated. The residue was purified via silica gel chromatography with ethyl acetate-hexane (10 to 60 %). Evaporation of the desired fractions resulted in 639 mg (74.8 %) of the title compound as an oil.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.60-7.10 (m, 11 H), 6.60  
10 (m, 2 H), 5.10 (s, 2 H), 4.75 (br d, 1 H), 3.80 (app d, 3 H), 3.55 (br d, 1 H), 3.00-2.50 (m, 4 H), 1.80 (m, 2 H), 1.60 (m, 1H), 1.30 (m, 1 H), 1.10 (m, 1 H). MS (ESI)  $m/e$  414 ( $m^+$ ).

### Example 11B

#### Preparation of 4-benzylpiperidin-1-yl-(2-methoxy-4-hydroxy)benzene carboxamide



15 A Par hydrogenation vessel was charged with 4-benzylpiperidin-1-yl-(2-methoxy-4-benzyloxy)benzene carboxamide from Example 11A (15.00 g, 36.14 mM), dissolved in MeOH (100 mL), and 5 % palladium on carbon (1.2 g). The vessel was flushed, placed under 42 psi of  $\text{H}_2$ , and shaken for 16 h. The slurry was then filtered through Celite and  
20 concentrated. The crude material was recrystallized from ethyl acetate/hexane (1 : 10) to yield 11.24 g (95.7 %) of the title compound as a white solid.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.40-6.80 (m, 6 H), 6.30 (m, 2 H), 4.75 (broad d, 2 H), 3.50 (app d, 3 H), 3.0-2.50 (m, 4 H), 1.75 (m, 2 H), 1.55 (m, 1 H), 1.20 (m, 1 H), 1.05 (m, 1 H). MS (ESI)  $m/e$  324 ( $m^+$ ).

Example 12Effect of Ortho-Substitution

Using the methods set forth in Example 10, 1-benzoyl-4-benzylpiperidine isomers, substituted on the benzoyl substituent at various positions, were tested for their ability to inhibit p38 $\alpha$  kinase at 15  $\mu$ M concentration. The results are shown in Table 3:

Table 3: % Inhibition of p38 $\alpha$ kinase at 15 $\mu$ M			
	<i>ortho</i>	<i>meta</i>	<i>para</i>
F	98	92	94
Cl	99	80	85
Br	99	74	96
methyl	98	90	97
CF <sub>3</sub>	98	49	66

As shown in Table 3, for any given substituent, the ortho isomer shows enhanced inhibition.

The IC<sub>50</sub> values, obtained as set forth in Example 10, were also determined for the ortho-, meta-, and para-methoxy substituted benzoyl forms of 1-benzoyl-4-benzylpiperidine. These results are shown in Table 4:

Table 4: IC <sub>50</sub> 's of methoxy analogs	
	IC <sub>50</sub> ( $\mu$ M)
<i>ortho</i>	0.287
<i>meta</i>	1.1
<i>Para</i>	0.605

It is clear from these results that substitution at the ortho position is preferred.

The molar refraction values for the ortho derivatives containing chloro, bromo and methoxy substituents were determined as a measure of the size of the substituent as described by Hansch, C. *et al.*, *J Med Chem* (1973) 16:1207-1216. A linear correlation between the MR values and percent inhibition of p38 $\alpha$  at 200 nM concentrations of the compounds was obtained. The smallest substituent (chloro) showed an MR value of about 6 and 52% inhibition at 200 nM; the compound with an intermediate MR value of about 8 (methoxy) provided 68% inhibition at this concentration, and the compound having the




largest substituent (bromo) showed an MR value of approximately 9 and 72% inhibition at this concentration. Thus, it appears that the ability to inhibit p38 $\alpha$  kinase is dependent on the size of the ortho substituent. While not intending to be bound by any theory, applicants believe that the bulkier substituents rotate the aroyl moiety out of the plane of the piperidine or piperazine ring.

5

U.S. Provisional Applications Serial Nos. 60/098,219 and 60/125,343 are relied on and incorporated herein by reference in their entirety.

Claims

1. A method to treat a condition mediated by p38 $\alpha$  kinase which method comprises administering to a subject in need of such treatment a compound of the formula:



and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof,

wherein

Z is N or CR<sup>1</sup>, R<sup>1</sup> is a noninterfering substituent,

each of X<sup>1</sup> and X<sup>2</sup> is a linker,

Ar<sup>1</sup> and Ar<sup>2</sup> are identical or different, and represent optionally substituted C1-C20 hydrocarbyl residues wherein at least one of Ar<sup>1</sup> and Ar<sup>2</sup> is an optionally substituted aryl group, with the proviso that when X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof, X<sup>1</sup> is CO or an isostere thereof, and Ar<sup>2</sup> is optionally substituted phenyl, Ar<sup>1</sup> is other than an optionally substituted indolyl, benzimidazolyl or benzotriazolyl substituent, and wherein said optionally substituted phenyl is not an optionally substituted indolyl, benzimidazolyl, or benzotriazolyl,

Y is a noninterfering substituent, wherein n is an integer from 0-4, and

wherein m is an integer from 0-4 and l is an integer from 0-3.

2. The method defined in claim 1, wherein the R<sup>1</sup> is selected from the group consisting of optionally substituted alkyl, alkoxy, aryl, arylalkyl, aryloxy, heteroaryl, amino, ureayl, carbamate, polyhaloalkoxy, halogen, acyl, carboxy, or hydroxy.

3. The method defined in claim 1, wherein Y is selected from the group consisting of H, optionally substituted alkyl, alkoxy, cycloalkyl, cycloheteroalkyl, aryl, cycloheteroalkyl, heteroaryl, halogen, alkylaminocarbonyl, arylaminocarbonyl,

heteroarylaminocarbonyl, acyl, carboxy, hydroxy, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, amino, and alkylamino.

4. The method defined in claim 1, wherein said linker is saturated or  
5 unsaturated alkylene optionally containing 1-4 carbonyl, 1-4 SO<sub>2</sub>, and/or 1-3 heteroatoms, including a linker which is itself CO, SO<sub>2</sub>, SO or is itself a heteroatom, and said alkylene or suitable heteroatom optionally substituted with a substituent selected from the group consisting of halo, alkyl, alkoxy, cycloalkyl, aryl, aryloxy, arylalkyl, haloalkyl, polyhaloalkyl, haloalkoxy, polyhaloalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl,  
10 heteroaryl, heteroarylalkyl, heteroaryloxy, arylthio, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl, alkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, acyl, aminocarbonyl, arylcarbonyl, heteroarylcarbonyl, cyano, carboxy, hydroxy, tetrazolyl, imidazole, oxazole, triazole, and -SOR wherein R is  
15 hydroxy, alkyl, aryl, alkoxy, aryloxy, cycloalkyl, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy or cycloheteroalkylalkoxy.

5. The method defined in claim 1, wherein  
when Z is N, X<sup>2</sup> is the same as X<sup>1</sup>, or X<sup>2</sup> is -O-, -O[C(Rx)<sub>2</sub>]<sub>r</sub>- wherein Rx is  
20 optionally substituted alkyl, alkenyl or alkynyl and r is an integer from 1 to 4, -NR<sup>2</sup>CO-, -SO<sub>2</sub>NR<sup>2</sup>-, -NR<sup>2</sup>-, -S-, -SO-, SO<sub>2</sub>, wherein R<sup>2</sup> is H, optionally substituted alkyl, cycloalkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, carboxy, or acyl; or  
wherein when Z is CR<sup>1</sup>, X<sup>2</sup> is X<sup>1</sup> or is -O[C(Rx)<sub>2</sub>]<sub>r</sub> wherein Rx is -SO<sub>2</sub>-.

25 6. The method defined in claim 4 wherein said alkylene group is C1-C4.

7. The method defined in claim 1, wherein Ar<sup>1</sup> or Ar<sup>2</sup> is an aryl group which is the residue of an aromatic hydrocarbon containing one or more rings optionally including one or more heteroatoms, selected from the group of O, N and S.

8. The method defined in claim 4,  
wherein said haloalkyl or polyhaloalkyl group is CF<sub>3</sub> or CF<sub>3</sub>CH<sub>2</sub>; and  
wherein said haloalkoxy or polyhaloalkoxy group is CF<sub>3</sub>O or CF<sub>3</sub>CH<sub>2</sub>O.

5 9. The method defined in claim 7, wherein said aryl group has 6 to 12 carbon  
atoms and 0 to 3 heteroatoms.

10 10. The method defined in claim 9, wherein said aryl group has 6 to 8 carbon  
atoms and 1 or 2 heteroatoms.

11 11. The method defined in claim 7, wherein said aryl group is phenyl or  
residues of optionally benzo-fused heterocycles containing up to 3 heteroatoms selected  
from the group consisting of S, N and O.

15 12. The method defined in claim 7, wherein said aryl group is saturated or  
unsaturated 5- to 7-membered heterocycle.

13. The method defined in claim 12, wherein said aryl group is saturated or  
unsaturated 5- to 6-membered heterocycle.


20 14. The method defined in claim 11, wherein said aryl group is selected from  
the group consisting of indolyl, isoquinolyl, quinolyl, benzimidazolyl, benzotriazolyl,  
benzothiazolyl, benzofuranyl, pyridyl, thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl,  
imidazolyl, morpholinyl or piperidyl.

25 15. The method defined in claim 1 wherein said linker is not CO or an isostere  
thereof, and

wherein at least one of said Ar<sup>1</sup> and Ar<sup>2</sup> is indolyl, benzimidazolyl or  
benzotriazolyl.

30

16. A method to treat a condition mediated by p38 $\alpha$  kinase defined in claim 1 which method comprises administering to a subject in need of such treatment a compound of the formula:



5 and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof,

wherein Ar<sup>1</sup> is optionally substituted furanyl, thiophenyl, phenyl system containing 0, 1 or 2 N as heterocyclic atoms, or naphthyl system containing 0, 1, 2 or 3 N as heterocyclic atoms,

10 X<sup>1</sup> is CO or an isostere thereof;

Y is optionally substituted alkyl (1-6C), optionally substituted aryl (6-10C), or optionally substituted arylalkyl (7-11C);

n is 0-1;

Z is N or CH;

15 X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof; and

Ph is optionally substituted phenyl.

17. The method of claim 16 wherein Ph is unsubstituted, or wherein n is 0, or

20 wherein Z is CH.

18. The method of claim 17 wherein n is 0 and Z is CH.

19. The method of any of claims 16-18 wherein Ar<sup>1</sup> is optionally substituted  
25 phenyl.

20. The method of claim 16 wherein Ar<sup>1</sup>X<sup>1</sup> is benzoyl, 4-methoxybenzoyl, 4-cyanobenzoyl, 4-phenylbenzoyl, 3-methoxybenzoyl, 3,5-dimethoxybenzoyl, 2-chlorobenzoyl, 2-bromobenzoyl, 2-iodobenzoyl, 2-methoxybenzoyl, 2-methylbenzoyl, 2-(2-carbomethoxy ethen-1-yl) benzoyl, 2-naphthoyl, 1-naphthoyl, 2-furanoyl, 2-thiophenoyl, 2-dimethylaminobenzoyl, 3-dimethylaminobenzoyl, 4-dimethylaminobenzoyl, 2-pyridoyl, 3-pyridoyl, 4-pyridoyl, 3,4,5-trimethoxybenzoyl, 2-acetamidobenzoyl, 3-acetamidobenzoyl, 4-acetamidobenzoyl, 4-formamidobenzoyl, 2-aminobenzoyl, 3-aminobenzoyl, 4-aminobenzoyl, 4,5-dimethoxybenzoyl, 2,4-dimethoxybenzoyl, 2-hydroxybenzoyl, 2-benzoyloxybenzoyl, 3,5-bis-trifluoromethylbenzoyl, 2,6-dimethoxybenzoyl, 2-methylaminobenzoyl, 3-methylaminobenzoyl, 4-methylaminobenzoyl, 2-hydroxy-4-aminobenzoyl, 2-chloro-4-aminobenzoyl, 2-chloro-4-methylaminobenzoyl, 2-chloro-4-dimethylaminobenzoyl, 2-methoxy-4-nitrobenzoyl, 2-methoxy-4-aminobenzoyl, 4-guanidinobenzoyl, phenylsulfonyl, 4-amino-3-pyridoyl, 2-aminobenzoyl, 2-methoxy-4-hydroxybenzoyl, 2-methoxy-4-benzoyloxybenzoyl, or 2-methoxy-4-methoxybenzoyl.

21. The method of claim 16 wherein formula (1) is 1-benzoyl-4-benzyl piperidine, 1-(3-methoxybenzoyl)-4-benzyl piperidine, 1-(4-phenylbenzoyl)-4-benzyl piperidine, 1-(2-chlorobenzoyl)-4-benzyl piperidine, 1-(2-bromobenzoyl)-4-benzyl piperidine, 1-(2-iodobenzoyl)-4-benzyl piperidine, 1-(2-methoxybenzoyl)-4-benzyl piperidine, 1-(2-methylbenzoyl)-4-benzyl piperidine, 1-(2-naphthoyl)-4-benzyl piperidine, 1-(4-dimethylaminobenzoyl)-4-benzyl piperidine, 1-(4-acetamidobenzoyl)-4-benzyl piperidine, 1-(4-formamidobenzoyl)-4-benzyl piperidine, 1-(4-aminobenzoyl)-4-benzyl piperidine, 1-(2,4-dimethoxybenzoyl)-4-benzyl piperidine, 1-(2,6-dimethoxybenzoyl)-4-benzyl piperidine, 1-(2-chloro-4-aminobenzoyl)-4-benzyl piperidine, 1-(2-chloro-4-methylaminobenzoyl)-4-benzyl piperidine, 1-(2-chloro-4-dimethylaminobenzoyl)-4-benzyl piperidine, *cis*-1-(2-bromobenzoyl)-3-methyl-4-benzyl piperidine, 1-(2-methoxy-4-hydroxybenzoyl)-4-benzyl piperidine, 1-(2-methoxy-4-methoxybenzoyl)-4-benzyl piperidine, 1-(2-methoxy-4-benzoyloxybenzoyl)-4-benzyl piperidine, or 1-(2-methoxy-4-methoxybenzoyl)-4-(4-fluorobenzyl)piperidine.

22. The method of claim 16 wherein said condition is characterized by a proinflammation response.





5           23. The method of claim 22 wherein said condition characterized by inflammation is acute respiratory distress syndrome, asthma, chronic obstructive pulmonary disease, uveitis, acute renal failure, head trauma, ischemic/reperfusion injury, multiple sclerosis, IBD, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, other arthritic conditions, sepsis, septic shock, endotoxic shock, Gram-negative  
10 sepsis, toxic shock syndrome, adult respiratory distress syndrome, stroke, reperfusion injury, CNS injury, psoriasis, restenosis, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcosis, a bone resorption disease, graft-versus-host reaction, Crohn's Disease, ulcerative colitis, or pyresis.

15           24. The method of claim 16 wherein said condition is a heart condition associated with cardiac failure.

          25. The method of claim 24 wherein said chronic heart condition is congestive heart failure, cardiomyopathy or myocarditis.


20

          26. The method defined in claim 1 wherein the compound is at least one compound selected from the following list:


Compound #	Structure
1	
2	
3	
4	




5




6




7




8




9




10




11




12




13




14




15




16




17




18




19




20




21




22




23




24
















25




26




27	
28	
29	
30	
31	
32	
33	

34	 <chem>COc1ccc(cc1)C(=O)N2Cc3ccccc3cc2C(=O)N2Cc4ccccc4</chem>
35	 <chem>COc1ccc(cc1)C(=O)N2Cc3ccccc3cc2C(=O)N2Cc4ccccc4</chem>
36	 <chem>CC1=C(C)N2C(=C(C1)C(=O)N2Cc3ccccc3)C4=CC=CC=C4</chem>
37	 <chem>c1cc2nc3ccccc3cc2s1C(=O)N2Cc3ccccc3cc2</chem>
38	 <chem>CC(C)Nc1cc(C(=O)N2Cc3ccccc3cc2)ccn1</chem>
39	 <chem>c1cc2nnc3ccccc3cc2c1C(=O)N2Cc3ccccc3cc2</chem>


40




41




42




43




44




45




46




47










48



















49






50	
51	
52	
53	
54	
55	
56	


57	
58	
59	
60	
61	
62	
63	
64	

65	
66	
67	
68	
69	
70	
71	
72	


73




74




75




76




77




78




79




80




81




82




83




84















85




86




87	
88	
89	
90	
91	
92	




99




100




101




102



103




104







105




106




107




108




109



110






27. The method defined in claim 1 further comprising administering said compound and an additional therapeutic agent.

5

28. A pharmaceutical composition for treating conditions characterized by enhanced p38- $\alpha$  activity which composition comprises  
a therapeutically effective amount of a compound of the formula



10 and the pharmaceutically acceptable salts thereof, or a pharmaceutical composition thereof,

wherein

Z is N or CR<sup>1</sup>, R<sup>1</sup> is a noninterfering substituent,

each of X<sup>1</sup> and X<sup>2</sup> is a linker,

15 Ar<sup>1</sup> and Ar<sup>2</sup> are identical or different, and represent optionally substituted C1-C20 hydrocarbyl residues wherein at least one of Ar<sup>1</sup> and Ar<sup>2</sup> is an optionally substituted aryl group, with the proviso that when X<sup>2</sup> is CH<sub>2</sub> or an isostere thereof, X<sup>1</sup> is CO or an isostere thereof, and Ar<sup>2</sup> is optionally substituted phenyl, Ar<sup>1</sup> is other than an optionally substituted

indolyl, benzimidazolyl or benzotriazolyl substituent, and wherein said optionally substituted phenyl is not an optionally substituted indolyl, benzimidazolyl, or benzotriazolyl,

5           Y is a noninterfering substituent, wherein n is an integer from 0-4, and  
          wherein m is an integer from 0-4 and l is an integer from 0-3.

29.     The composition of claim 28 which further contains an additional therapeutic agent.

10           30.     The composition of claim 29 wherein said additional therapeutic agent is a corticosteroid, a monoclonal antibody, or an inhibitor of cell division.